

PAIN INTERFERENCE

A brief guide to the PROMIS Pain Interference instruments:

ADULT	ADULT CANCER	PEDIATRIC	PARENT PROXY
PROMIS Item Bank v1.0 – Pain Interference	PROMIS-Ca Bank v1.0 – Pain Interference	PROMIS Pediatric Bank v1.0 – Pain Interference	PROMIS Parent Proxy Bank v1.0 – Pain Interference
PROMIS Item Bank v1.1 – Pain Interference			
PROMIS Short Form v1.0 – Pain Interference 4a	PROMIS-Ca Bank v1.1 – Pain Interference	PROMIS Pediatric Short Form v1.0 – Pain Interference 8a	PROMIS Parent Proxy Short Form v1.0 – Pain Interference 8a
PROMIS Short Form v1.0 – Pain Interference 6a			
PROMIS Short Form v1.0 – Pain Interference 6b			
PROMIS Short Form v1.0 – Pain Interference 8a			

ABOUT PAIN INTERFERENCE

The PROMIS Pain Interference instruments measure the self-reported consequences of pain on relevant aspects of one’s life. This includes the extent to which pain hinders engagement with social, cognitive, emotional, physical, and recreational activities. Pain Interference also incorporates items probing sleep and enjoyment in life, though the item bank only contains one sleep item. The Pain Interference short form is universal rather than disease-specific. It assesses pain interference over the past seven days.

Pain Interference instruments are available for adults (ages 18+), pediatric self-report (ages 8-17) and for parents serving as proxy reporters for their child (youth ages 5-17).

(For complete definition see <http://nihpromis.org/measures/domainframework1>)

INTRODUCTION TO ASSESSMENT OPTIONS

There are two administration options for assessing Pain Interference: short forms and computerized adaptive test (CAT). When administering a short form, instruct participants to answer all of the items (i.e., questions or statements) presented. With CAT, participant responses guide the system’s choice of subsequent items from the full item bank (40 items in total). Although items differ across respondents taking CAT, scores are comparable across participants. Some administrators may prefer to ask the same question of all respondents or of the same respondent over time, to enable a more direct comparability across people or time. In these cases, or when paper administration is preferred, a short form would be more desirable than CAT. This guide provides information on all Pain Interference short form and CAT instruments.

Whether one uses a short form or CAT, the score metric is Item Response Theory (IRT), a family of statistical models that link individual questions to a presumed underlying trait or concept of pain interference represented by all items in the item bank. When choosing between CAT and a short form, it is useful to consider the demands of computer-based assessment, and the psychological, physical, and cognitive burden placed on respondents as a result of the number of questions asked.




Figure 1

Figure 1 illustrates the correlations (strength of relationship) of the full bank with CAT and with short

forms of varying length. The correlation of CAT scores with the full bank score is greater than a short form of any length. A longer CAT or longer short form offers greater correlation, as well as greater precision. When evaluating precision, not all questions are equally informative. The flexibility of CAT to choose more informative questions offers more precision.

Some PROMIS domains have multiple versions of instruments (i.e. Item Banks/Computerized Adaptive Tests (CATs) and Short Forms). Generally, it is recommended that you use the most recent version available which can be identified as the instruments with the highest version number.

Instruments are changed for various reasons. For example, the original PROMIS Bank version 1.0 – Physical Function included 124 items after being tested in a diverse sample. Later, during an effort to translate instrument items into multiple languages, translation challenges were identified. Therefore, minor modifications to the English source items were required (e.g. metric equivalents to measurements such as “Over 10 pounds/ 5 kg” were added). These modifications (19 in total) resulted in the creation of a version 1.1 item bank. Later, version 1.2 was created by eliminating two items due to restrictions in their use.

In most cases, an instrument that has a decimal increase (v1.0 to v1.1 or v1.2) retains the same item-level parameters as well as instrument reliability and validity. In cases where a version number increases by a whole number (e.g., v1.0 to v2.0), the changes to the instrument are more substantial. For example, the PROMIS Bank v1.0 – Satisfaction with Participation in Social Roles is a small item bank comprised of 14 role function items. While the PROMIS Bank v2.0 – Satisfaction with Social Roles and Activities instrument includes a broader range of item content. Version 2.0 was re-calibrated independent of version 1.0, and is a larger (44 items), superior item bank. Details on the specific differences between instrument versions can be found in scoring manuals available on the Assessment Center homepage.

SHORT FORM DIFFERENCES

You will notice that there are 4 Pain Interference short forms for adults. Items in the 4a, 6a, and 8a short forms were selected based on rankings using two psychometric criteria: (1) maximum interval information; and 2) CAT simulations. Item rankings were similar for both criteria. For the maximum interval criterion, each item information function was integrated (without weighting) for the interval from the mean to 2 SDs worse than the mean. For the CAT simulations, responses to all items in each bank were generated using a random sample of 1,000 simulees drawn separately for each bank (centered on 0.5 SD worse than the general population mean). Items were rank ordered based on their average administration rank over the simulees. Content experts reviewed the items and rankings and made cuts of 4, 6, and 8 items. For each domain, 4-item, 6-item and 8-items have been selected so that the items are nested/overlap (e.g., the 8-item form is the 6-item form plus two additional items). The 4a, 6a, and 8a short forms can be administered with short forms of similar length from other domains (Depression, Physical Function, Anxiety, Fatigue, Sleep Disturbance, and Satisfaction with Participation in Social Roles v1.0) as part of a PROMIS Profile (see PROMIS-29, 43 or 57 Profile v2.0), though they can also be administered individually.

The original short form (6b) was constructed by the domain team with a focus on representing the range of the trait and also representing the content of the item bank. Domain experts reviewed short forms to give input on the relevance of each item. Each domain group worked independently and the original short forms are 6-10

items long depending on the domain. Psychometric properties and clinical input were both used and likely varied in importance across domains.

In selecting between short forms, the difference is instrument length. The reliability and precision of the short forms within a domain is highly similar. If you are working with an adult sample in which you wanted the most precise measure, select the 8a short form. If you are working in an adult sample in which you expected huge variability in a domain area and wanted different subdomains covered, you should select the 6b short form. If you had little room for additional measures but really wanted to capture something as a secondary outcome, you should use one of the shorter instruments (4a, 6a). For pediatric self-report and parent proxy report, there is only one short form available.

SELECTING THE ADULT CANCER INSTRUMENT

In selecting whether to use the adult cancer instrument (PROMIS-Ca) for this domain, it is important to consider the patient population being studied. All PROMIS-Ca instruments were developed for use with any cancer patient. This was done by having content experts review the adult PROMIS item bank to determine if there was a need to develop additional items or remove items because they conveyed a different meaning in cancer. Next, calibration testing with cancer patients with different diagnoses and treatments was conducted and data was analyzed to determine the final set of items and calibrations. The PROMIS-Ca Pain Interference CAT contains a total of 35 items, 32 of which are also in the PROMIS Pain Interference CAT. Some PROMIS-Ca items use calibrations that are different from the PROMIS adult calibrations. The Cancer calibration sample is selected by default when you add the PROMIS-Ca CATs to a study in Assessment Center. There are no short form instruments created specifically for an adult cancer population in Assessment Center.

SELECTING A PEDIATRIC OR PARENT PROXY INSTRUMENT

In selecting whether to use the pediatric or parent proxy instrument for this domain, it is important to consider both the population and the domain which you are studying. Pediatric self-report should be considered the standard for measuring patient-reported outcomes among children. However, circumstances exist when the child is too young, cognitively impaired, or too ill to complete a patient-reported outcome instrument. While information derived from self-report and proxy-report is not equivalent, it is optimal to assess both the child and the parent since their perspectives may be independently related to healthcare utilization, risk factors, and quality of care.

WHICH CALIBRATION SAMPLE SHOULD I USE?

The PROMIS Parent Proxy instruments have two calibration samples – Parent Proxy and Parent Proxy without Local Dependence. The former includes calibrations for all items. This is the default calibration sample. If you aren't sure which calibration sample to use, utilize this one. The Parent Proxy without Local Dependence does not include calibrations for some items. The items without calibrations are enemy items. That is, a dyad or triad of items was identified in which there are psychometric reasons to only administer one of those items to a given respondent. For example, item Pf3pain4 and Pf3pain2 are enemy items. A participant should only see one of these items in a CAT.



SCORING THE INSTRUMENT

Short Forms: PROMIS instruments are scored using item-level calibrations. This means that the most accurate way to score a PROMIS instrument is to utilize scoring tools within Assessment Center or API that look at responses to each item for each participant. Data collected in either of these platforms will automatically score in this way. We refer to this as “response pattern scoring.” Response pattern scoring can be used when data was collected on paper or in another software package through the [Assessment Center Scoring Service](#). Because response pattern scoring is more accurate than the use of raw score/scale score look up tables, it is preferred. However, if you aren’t able to use response pattern scoring, you can use the instructions below which rely on raw score/scale score look-up tables.

For adults, each question has five response options ranging in value from one to five (for pediatrics and parent proxy it is 0 to 4). To find the total raw score for a short form with all questions answered, sum the values of the response to each question. For example, for the adult 6-item form, the lowest possible raw score is 6; the highest possible raw score is 30 (see all short form scoring tables in Appendix 1).

A score can be approximated if a participant skips a question. If items are missing, first check how many items were answered. For short forms with at least 5 items, confirm that 4 or 50% of items, whichever is greater, were answered. For example, a 4-item short form can only be scored with complete data. A 5-item short form can be scored as long as 4 items were answered. A 10-item short form can be scored as long as the participant answered at least 5 items. For branched instruments (e.g., Alcohol Use), the screening question is not used in calculating the score and therefore shouldn’t be counted when assessing if the minimum number of items were answered. After confirming that enough responses were provided, sum the response scores from the items that were answered (not including any screening question). Multiply this sum by the total number of items in the short form. Finally, divide by the number of items that were answered. For example, if a respondent answered 5 of 8 questions and answered all items with the second lowest response option (2), you would sum all responses (10), multiply by the number of items in the short form (8) and divide by the number of items that were answered (5). Here $(10 \times 8) / 5 = 16$. If the result is a fraction, round up to the nearest whole number. This is a pro-rated raw score.

Again, the formula is:

$$\frac{(\text{Raw sum} \times \text{number of items on the short form})}{\text{Number of items that were actually answered}}$$

Locate the applicable score conversion table in Appendix 1 and use this table to translate the total raw score or pro-rated score into a T-score for each participant. The T-score rescales the raw score into a standardized score with a mean of 50 and a standard deviation (SD) of 10. Therefore a person with a T-score of 40 is one SD below the mean. It is important to note that Assessment Center will convert a participant’s pattern of responses to a standardized T-score after they have finished a CAT. The standardized T-score is reported as the final score for each participant.

For the adult PROMIS Pain Interference 6b short form, a raw score of 10 converts to a T-score of 53.8 with a standard error (SE) of 1.8 (see scoring table for the 6b short form in Appendix 1). Thus, the 95% confidence interval around the observed score ranges from 50.3 to 57.3 (T-score $\pm (1.96 \times SE)$ or $53.8 \pm (1.96 \times 1.8)$).

For pro-rated scores, this calculation assumes that responses are missing at random. This isn’t always true. Therefore, use caution when interpreting the final pro-rated T-score.

CAT: A minimum number of items (4 for adult and adult cancer CATs and 5 for Peds and Parent Proxy CATs) must be answered in order to receive a score for Pain Interference CAT. The first item is selected because it provides the most information about the U.S. general population (or a general cancer population in the case of the adult cancer instruments). The response to this item will guide the system’s choice of the next item for the participant. The participant’s response to this item will dictate the selection of the following question, and so on. As additional items are administered, the potential for error is reduced and confidence in the respondent’s score increases. CAT will continue until either the standard error drops below a specified level, or the participant has answered the maximum number of questions (12), whichever occurs first.

For most PROMIS instruments, a score of 50 is the average for the United States general population with a standard deviation of 10 because calibration testing was performed on a large sample of the general population. However, adult cancer, pediatric and parent proxy instruments were not calibrated on a national sample and so a score of 50 represents the average of the calibration sample which was generally more enriched for chronic illness (i.e. cancer). In these instruments, a score of 50 likely represents somewhat sicker people than the general population. The T-score is provided with an error term (Standard Error or SE). The Standard Error is a statistical measure of variance and represents the “margin of error” for the T-score.

Important: A higher PROMIS T-score represents more of the concept being measured. For negatively-worded concepts like Pain Interference, a T-score of 60 is one SD worse than average. By comparison, a Pain Interference T-score of 40 is one SD better than average.

STATISTICAL CHARACTERISTICS

There are four key features of the score for Pain Interference:

- **Reliability:** The degree to which a measure is free of error. It can be estimated by the internal consistency of the responses to the measure, or by correlating total scores on the measure from two time points when there has been no true change in what is being measured (for z-scores, reliability = $1 - SE^2$).
- **Precision:** The consistency of the estimated score (reciprocal of error variance).
- **Information:** The precision of an item or multiple items at different levels of the underlying continuum (for z-scores, information = $1/SE^2$).
- **Standard Error (SE):** The possible range of the actual final score based upon the scaled T-score. For example, with a T-score of 52 and a SE of 2, the 95% confidence interval around the actual final score ranges from 48.1 to 55.9 (T-score \pm (1.96*SE) = $52 \pm 3.9 = 48.1$ to 55.9).

The final score is represented by the T-score, a standardized score with a mean of 50 and a standard deviation (SD) of 10.




Figure 2




Figure 3

In Figure 2 (adult 6b short form), the two dotted horizontal lines each represent a degree of internal consistency reliability (i.e., .90 or .95) typically regarded as sufficient for an accurate individual score. The shaded blue region marks the range of the scale where measurement precision is comparable to the reliability of .90 for the six-item form. Figure 2 also tells us where on the scale the form is most informative based upon the T-score. This form would typically be more informative than a Pain Interference form with fewer items.

Scaling Model Used For Calibration	Promis	
Total Number of Items	41	
Sample	N	Alpha Reliability
Promis	15903	0.99

Score Distributions											
	Mean	SD	P5	P10	P25	P50	P75	P90	P95		
Raw	87.14	51.51	33.00	36.00	42.00	66.00	131.00	170.00	183.00		
Scale	55.92	10.82	39.98	39.98	47.48	55.22	65.34	70.30	72.84		
Conditional Reliability											
Scale Score	10.0	20.0	30.0	40.0	50.0	60.0	70.0	80.0	90.0	Min	Max
SE		51.60	12.60	2.80	.50	.10	.10	.10	.20	.70	
Reliability		.00	.00	.00	.71	.99	1.00	.99	.96	.48	

Figure 4

Figure 3 (Adult 4a, 6a & 8a short forms) also tells us where on the scale the form is most informative based upon the T-score: the 8-item form is more informative than the 6-item form, which is more informative than the 4-item form. See additional test information figures for Pediatric and Parent Proxy instruments in Appendix 2.

Figure 4 is a sample of the statistical information available in Assessment Center for the adult Pain Interference CAT.

More information is available online via Assessment Center (assessmentcenter.net).

PREVIEW OF SAMPLE ITEM

Figure 5 shows a Pain Interference item from the adult full item bank as it would appear to a study participant during data collection in Assessment Center. Several formats for presenting the items are available for computer-based administration through Assessment Center (see FAQ section).




Figure 5

In the past 7 days...		Not at all	A little bit	Somewhat	Quite a bit	Very much
PAININ9 1	How much did pain interfere with your day to day activities?	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5
PAININ22 2	How much did pain interfere with work around the home?	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5
PAININ31 3	How much did pain interfere with your ability to participate in social activities?	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5

Figure 6

Figure 6 is an excerpt from the paper version of the adult six-item short form. This is the paper version format used for all Pain Interference instruments. It is important to note, CAT is not available for paper administration.

DATA REPORTS

Upon completion of an adult CAT for Physical Function, Depression, Anxiety, Pain Interference, Fatigue, Sleep Disturbance, and Satisfaction with Participation in Social Roles v1.0, a data report is available in Assessment Center. Figure 7 demonstrates some of the information available on the data reports.




Figure 7

To access a sample report for Pain Interference, complete the CAT demo at nihpromis.org. More than one CAT domain can be completed at a time; results for all domains selected will be generated and displayed within the one report.

Data reports are also available if you choose to administer an adult PROMIS Profile instrument, which includes a short form from seven PROMIS domains (Physical Function, Depression, Anxiety, Fatigue, Pain Interference, Satisfaction with Participation in Social Roles, and Sleep Disturbance).

FREQUENTLY ASKED QUESTIONS (FAQ)

Q: I am interested in learning more. Where can I do that?

All instruments are available on the PROMIS website through Assessment Center, which houses all PROMIS instruments for each domain.

Assessment Center is a free online research management tool. It enables researchers to create study-specific websites for capturing participant data securely. Studies can include measures within the Assessment Center library, as well as custom instruments created or entered by the researcher. PROMIS instruments (short forms, CATs, profiles) are a central feature of the instrument library within Assessment Center. Any PROMIS measure can be included in an online study or downloaded for administration on paper.

Detailed statistical information and development history about PROMIS items and instruments are available for review at nihpromis.org or assessmentcenter.net. To learn more, contact help@assessmentcenter.net.



Q: Do I need to register with PROMIS to use these instruments?

Yes, to get a copy of these instruments, we ask that you register with Assessment Center and endorse the PROMIS terms and conditions of use, so that we are better able to track who has accessed instruments for research. Assessment Center is available at assessmentcenter.net.

Q: Are these instruments available in other languages?

Yes, these instruments are currently available in Spanish and German in Assessment Center. The PROMIS group is also working to translate this form into other languages. Information on available translations is updated periodically at <http://nihpromis.org/measures/translations>.

Q: Can I make my own short form?

Yes, custom Pain Interference short forms can be made by selecting any items from the item bank. Instructions for creating a custom short form in Assessment Center can be found in the Assessment Center User Manual <https://www.assessmentcenter.net/UserManuals.aspx>.

Q: How do I handle multiple responses when administering a short form on paper?

Guidelines on how to deal with multiple responses have been established. Resolution depends on the responses noted by the research participant.

- If two or more responses are marked by the respondent, and they are next to one another, then a data entry specialist will be responsible for randomly selecting one of them to be entered and will write down on the form which answer was selected. *Note: To randomly select one of two responses, the data entry specialist will flip a coin (heads - higher number will be entered; tails – lower number will be entered). To randomly select one of three (or more) responses, a table of random numbers should be used with a statistician’s assistance.*
- If two or more responses are marked, and they are NOT all next to one another, the response will be considered missing.

Q: What is the minimum change on a PROMIS instrument that represents a clinically meaningful difference?

This question is related to an area of active research in the PROMIS network, namely the determination of the “minimally important difference” or “MID” for a PROMIS instrument. A manuscript in the *Journal of Clinical Epidemiology* outlines the process for MIDs for adult PROMIS measures and estimates the MIDs for six PROMIS-Cancer scales: Yost, K. J., Eton, D. T., Garcia, S. F., & Cella, D. (2011). Minimally important differences were estimated for six PROMIS-Cancer scales in advanced-stage cancer patients. *Journal of Clinical Epidemiology*, 64(5), 507-16.

As described in that manuscript, the MID is a tool to enhance the interpretability of patient-reported outcomes and is often defined as the “the smallest difference in score in the domain of interest which patients perceive as beneficial and which would mandate, in the absence of troublesome side effects and excessive cost, a change in the patient’s management” (Jaeschke R, Singer J, Guyatt GH. Measurement of health status. Ascertaining the minimal clinically important difference. *Controlled Clinical Trials* 1989; 10(4):407-415).



APPENDIX 1-SCORING TABLES

Pain Interference 6b Short Form Conversion Table		
Raw Score	T-score	SE*
6	41.0	6.0
7	48.5	2.6
8	50.8	2.1
9	52.5	1.9
10	53.8	1.8
11	55.0	1.7
12	56.1	1.7
13	57.1	1.7
14	58.1	1.7
15	59.1	1.6
16	60.0	1.6
17	60.9	1.6
18	61.8	1.6
19	62.7	1.6
20	63.6	1.6
21	64.5	1.6
22	65.5	1.6
23	66.4	1.6
24	67.4	1.7
25	68.5	1.7
26	69.6	1.8
27	70.9	1.9
28	72.4	2.1
29	74.4	2.4
30	78.3	3.7

*SE = Standard Error on T-score metric

Adult version

Pain Interference 4a Short Form Conversion Table		
Raw Score	T-score	SE*
4	41.6	6.1
5	49.6	2.5
6	52.0	2.0
7	53.9	1.9
8	55.6	1.9
9	57.1	1.9
10	58.5	1.8
11	59.9	1.8
12	61.2	1.8
13	62.5	1.8
14	63.8	1.8
15	65.2	1.8
16	66.6	1.8
17	68.0	1.8
18	69.7	1.9
19	71.6	2.1
20	75.6	3.7

*SE = Standard Error

Pain Interference 6a Short Form Conversion Table		
Raw Score	T-score	SE*
6	41.1	6.0
7	48.6	2.4
8	50.7	1.8
9	52.2	1.6
10	53.4	1.6
11	54.5	1.6
12	55.6	1.5
13	56.6	1.5
14	57.6	1.5
15	58.6	1.5
16	59.5	1.5
17	60.4	1.4
18	61.2	1.4
19	62.1	1.4
20	63.0	1.5
21	63.8	1.5
22	64.8	1.5
23	65.7	1.5
24	66.7	1.5
25	67.6	1.5
26	68.7	1.5
27	69.8	1.5
28	71.0	1.6
29	72.6	2.0
30	76.3	3.6

*SE = Standard Error

Pain Interference 8a Short Form Conversion Table		
Raw Score	T-score	SE*
8	40.7	5.9
9	47.9	2.4
10	49.9	1.8
11	51.2	1.5
12	52.3	1.4
13	53.2	1.4
14	54.1	1.4
15	55.0	1.4
16	55.8	1.4
17	56.6	1.4
18	57.4	1.3
19	58.1	1.3
20	58.8	1.3
21	59.5	1.3
22	60.2	1.3
23	60.8	1.3
24	61.5	1.3
25	62.1	1.3
26	62.8	1.3
27	63.5	1.3
28	64.1	1.3
29	64.8	1.3
30	65.5	1.3
31	66.2	1.3
32	66.9	1.3
33	67.7	1.3
34	68.4	1.3
35	69.2	1.3
36	70.1	1.4
37	71.0	1.4
38	72.1	1.6
39	73.5	2.0
40	77.0	3.5

*SE = Standard Error

Adult version



Pain Interference 8a		
<i>Short Form Conversion Table</i>		
Raw Score	T-score	SE*
0	34.0	5.6
1	38.7	4.4
2	40.6	4.2
3	42.7	3.8
4	44.3	3.7
5	45.8	3.4
6	47.1	3.3
7	48.4	3.2
8	49.5	3.2
9	50.6	3.1
10	51.7	3.1
11	52.7	3.1
12	53.7	3.0
13	54.7	3.0
14	55.7	3.0
15	56.6	3.0
16	57.6	3.0
17	58.5	3.0
18	59.5	3.0
19	60.4	3.0
20	61.4	3.0
21	62.4	3.0
22	63.4	3.0
23	64.4	3.0
24	65.4	3.1
25	66.5	3.1
26	67.6	3.2
27	68.8	3.2
28	70.1	3.3
29	71.5	3.4
30	73.2	3.7
31	75.0	3.8
32	78.0	4.3

* SE = Standard error

Pediatric version

Parent Proxy Pain Interference		
<i>8-item Short Form Conversion Table</i>		
Summed Raw Score	Standard T-Score	SE*
0	38	6
1	44	3
2	46	3
3	48	3
4	49	2
5	50	2
6	51	2
7	52	2
8	53	2
9	54	2
10	55	2
11	56	2
12	57	2
13	58	2
14	58	2
15	59	2
16	60	2
17	61	2
18	62	2
19	62	2
20	63	2
21	64	2
22	65	2
23	66	2
24	67	2
25	67	2
26	68	2
27	69	2
28	70	2
29	71	3
30	73	3
31	74	3
32	78	4

*SE = Standard Error on T-score metric
Parent Proxy version

APPENDIX 2 – ADDITIONAL FIGURES




Figure 8 – Pediatric Test Information Pain